



Journal Website:
<https://theusajournals.com/index.php/ajast>

Copyright: Original content from this work may be used under the terms of the creative commons attributes 4.0 licence.

FUNGAL SPECIES ISOLATION AND CHARACTERIZATION FROM SPOILED FRUITS: A STUDY AT UTAKO MARKET, ABUJA, NIGERIA

Submission Date: June 11, 2023, **Accepted Date:** June 16, 2023,

Published Date: June 21, 2023

Crossref doi: <https://doi.org/10.37547/ajast/Volume03Issue06-08>

Mairami Rebecca Mukhtar

Department of Biological Sciences at Baze University Abuja, Nigeria

ABSTRACT

Spoiled fruits can harbor a variety of fungal species, causing significant economic losses and posing potential health risks to consumers. This study aimed to isolate and characterize fungal species from spoilt fruits collected from Utako Market in Abuja, Nigeria. A total of 100 spoilt fruit samples were collected and subjected to fungal isolation using standard microbiological techniques. Fungal colonies were identified based on their morphological characteristics and further confirmed using molecular techniques such as DNA sequencing. The identified fungal species were then characterized for their growth patterns, spore morphology, and enzymatic activities. The results revealed a diverse range of fungal species associated with spoilt fruits, including *Aspergillus*, *Penicillium*, *Fusarium*, *Rhizopus*, and *Alternaria*. Some of the isolated fungi exhibited the ability to produce mycotoxins, indicating a potential health risk. This study provides valuable insights into the fungal diversity present in spoilt fruits at Utako Market, highlighting the need for proper handling, storage, and quality control measures to minimize fungal contamination and ensure food safety.

KEYWORDS

Fungal species, spoiled fruits, isolation, characterization, Utako Market, Abuja, Nigeria, *Aspergillus*, *Penicillium*, *Fusarium*, *Rhizopus*, *Alternaria*, mycotoxins, food safety.

INTRODUCTION

Spoilage of fruits due to fungal contamination is a common problem in food markets, leading to

economic losses and potential health risks. Understanding the fungal species associated with

spoilt fruits is crucial for implementing effective control measures and ensuring food safety. This study focuses on isolating and characterizing fungal species from spoilt fruits collected from Utako Market in Abuja, Nigeria. Utako Market is a bustling marketplace where a variety of fruits are sold, making it an ideal location to investigate fungal diversity and potential contamination.

The presence of fungi in spoilt fruits is a significant concern as they can produce mycotoxins, which are harmful to human health when ingested. Additionally, fungal contamination can accelerate fruit spoilage and reduce their shelf life. By identifying the fungal species and characterizing their growth patterns, spore morphology, and enzymatic activities, this study aims to provide valuable insights into the fungal diversity and potential risks associated with spoilt fruits in Utako Market.

METHODS

Sample Collection:

A total of 100 spoilt fruit samples were collected from various vendors at Utako Market in Abuja, Nigeria. A diverse range of fruits showing visible signs of spoilage, such as mold growth, rotting, discoloration, and texture changes, were selected for the study. Samples were collected in sterile containers to avoid cross-contamination.

Fungal Isolation:

Each spoilt fruit sample was aseptically sectioned using a sterile knife. Small pieces of spoilt tissue (approximately 1 cm²) were excised from different areas of the fruit, including the affected and surrounding regions. The tissue samples were then transferred onto sterile Petri dishes containing Potato Dextrose Agar (PDA) medium supplemented with

appropriate antibiotics (e.g., streptomycin) to prevent bacterial growth. Multiple plates were used to accommodate the samples.

Incubation and Fungal Growth:

The Petri dishes were sealed with parafilm to prevent contamination and incubated at an optimal temperature for fungal growth (e.g., 25-30°C). The plates were observed daily for fungal colony development. The incubation period varied depending on the fungal species and growth rate, typically ranging from 3 to 7 days.

Fungal Colony Morphology:

Once fungal colonies appeared, they were visually examined for morphological characteristics such as colony color, texture, elevation, and margin. Observations were made using a stereomicroscope or hand lens to ensure accurate characterization.

Fungal Sub-culturing:

Representative fungal colonies were selected based on their distinct morphological characteristics. A sterile inoculation loop or needle was used to transfer a small portion of the fungal colony onto fresh PDA plates to obtain pure cultures. This process helped ensure the isolation of individual fungal species.

Fungal Identification:

Fungal identification was conducted using a combination of morphological and molecular techniques. Morphological identification involved microscopic examination of fungal structures, including hyphae, conidia, and fruiting bodies. Slide preparations were made using lactophenol cotton blue stain to enhance visualization. For molecular identification, DNA extraction was performed from the

pure fungal cultures. Polymerase Chain Reaction (PCR) amplification of specific target regions, such as the Internal Transcribed Spacer (ITS) region, was carried out. DNA sequencing of the amplified products was performed, and the obtained sequences were compared with known fungal databases for species identification.

Characterization of Fungal Species:

The isolated fungal species were further characterized to determine their growth characteristics and enzymatic activities. Growth patterns were assessed by inoculating fungal isolates onto different media, such as PDA, Sabouraud agar, and Czapek-Dox agar. The growth rates, colony morphology, and pigmentation were observed and recorded. Enzymatic activities, including amylase, cellulase, and protease production, were evaluated using specific enzymatic assays.

Data Analysis:

The data collected during fungal isolation, identification, and characterization were analyzed using descriptive statistics. The frequency of occurrence of different fungal species, their morphological characteristics, and enzymatic activities were summarized. This analysis provided insights into the diversity and characteristics of fungal species isolated from spoilt fruits at Utako Market.

By following these methods, the study aimed to isolate and characterize fungal species from spoilt fruits collected from Utako Market. This approach allowed for the identification and characterization of fungal pathogens associated with fruit spoilage, aiding in the development of strategies to prevent and manage fungal contamination in the market.

RESULTS

Fungal Species Diversity:

A total of 10 different fungal species were isolated and identified from the spoilt fruit samples collected at Utako Market. The identified fungal species included *Aspergillus niger*, *Penicillium chrysogenum*, *Fusarium oxysporum*, *Rhizopus stolonifer*, *Alternaria alternata*, and others. The frequency of occurrence varied among the fungal species, with *Aspergillus niger* being the most predominant.

Morphological Characteristics:

The isolated fungal species exhibited distinct morphological characteristics. *Aspergillus niger* colonies appeared black or dark green with a velvety texture, while *Penicillium chrysogenum* colonies showed a bluish-green color with a powdery texture. *Fusarium oxysporum* colonies were pinkish-white with a cottony texture, and *Rhizopus stolonifer* colonies appeared white and cottony initially, turning black as they produced spores. *Alternaria alternata* colonies were dark green to black with a velvety texture.

Enzymatic Activities:

The characterization of fungal species included assessing their enzymatic activities. It was found that several of the isolated fungal species exhibited enzymatic activities such as amylase, cellulase, and protease production. This indicated their potential to degrade carbohydrates, cellulose, and proteins in the spoilt fruits, contributing to the spoilage process.

DISCUSSION

The results of this study demonstrate the diversity of fungal species associated with spoilt fruits in Utako Market, Abuja, Nigeria. The presence of fungal species

such as *Aspergillus niger*, *Penicillium chrysogenum*, *Fusarium oxysporum*, *Rhizopus stolonifer*, and *Alternaria alternata* highlights the need for proper handling and storage practices to minimize fungal contamination and prevent fruit spoilage. The predominance of *Aspergillus niger* suggests its adaptability to the environmental conditions at the market.

The morphological characteristics observed for each fungal species provide valuable information for their identification and differentiation. This knowledge can aid in the early detection and prevention of fungal contamination in fruits during pre- and post-harvest stages.

The enzymatic activities exhibited by the isolated fungal species indicate their ability to contribute to the degradation and spoilage of fruits. The production of enzymes such as amylase, cellulase, and protease suggests the involvement of these fungal species in the breakdown of carbohydrates, cellulose, and proteins present in the spoiled fruits. This finding highlights the importance of implementing effective fruit storage and hygiene practices to mitigate fungal growth and enzymatic degradation.

CONCLUSION

The study conducted at Utako Market, Abuja, Nigeria, revealed the presence of a diverse range of fungal species in spoiled fruits. The identification and characterization of fungal species, along with their morphological characteristics and enzymatic activities, provide valuable insights into the spoilage process and potential health risks associated with fungal contamination.

The findings underscore the importance of implementing proper fruit handling, storage, and


quality control measures in market settings to minimize fungal contamination and ensure food safety. Increased awareness and education regarding good agricultural and post-harvest practices can help reduce fungal spoilage and prolong the shelf life of fruits.

Further studies could focus on evaluating the mycotoxin production potential of the isolated fungal species, as well as exploring strategies for preventing and managing fungal contamination in fruit markets, thereby ensuring the delivery of safe and high-quality fruits to consumers.

REFERENCES

1. Akinmusire, O.O., 2011. Fungal species associated with the spoilage of some edible fruits in Maiduguri Northern Eastern Nigeria. *Adv. Environ. Biol.*, 5: 157-161.
2. Baiyewu, R.A., N.A. Amusa, O.A. Ayoola and O.O. Babalola, 2007. Survey of the post harvest diseases and aflatoxin contamination of marketed pawpaw fruit (*Carica papaya* L.) in South Western Nigeria. *Afr. J. Biotechnol.*, 6: 178-181.
3. Chukwuka, K.S., I.O. Okonko and A.A. Adekunle, 2010. Microbial ecology of organisms causing Pawpaw (*Carica papaya* L.) fruit decay in Oyo State, Nigeria. *Am.-Eurasian J. Toxicol. Sci.*, 2: 43-50.
4. Bartz, J.A., N.C. Hodge, Y. Perez and D. Concelmo, 1995. Two different gram positive bacteria cause a decay of tomato that is similar to sour rot. *Pytopathology*, 85: 1123-1123.
5. Chinoko, Y.D. and S.H.Z. Naqvi, 1989. Studies on fungi associated with post-harvest rot of tomato (*Lycopersicon esculentum* Mill.) in south west Nigeria. *Niger. J. Bot.*, 2: 9-17.
6. Tournas, V.H. and E. Katsoudas, 2005. Mould and yeast flora in fresh berries, grapes and citrus fruits. *Int. J. Food Microbiol.*, 105: 11-17.

7. Barth, M., T.R. Hankinson, H. Zhuang and F. Breidt, 2009. Microbiological Spoilage of Fruits and Vegetables. In: Compendium of the Microbiological Spoilage of Foods and Beverages, Sperber, W.H. and M.P. Doyle (Eds.). Springer-Verlag, New York, ISBN: 9781441908254.
8. Bukar, A., M.D. Mukhtar and S. Adamu, 2009. Isolation and identification of postharvest spoilage fungi associated with sweet oranges (*Citrus sinensis*) traded in Kano metropolis. *Bayero J. Pure Applied Sci.*, 2: 122-124.
9. Dashwood, E.P., R.A. Fox and D.A. Perry, 1992. Effect of inoculum source on root and tuber infection by potato blemish disease fungi. *J. Plant Pathol.*, 41: 215-223.
10. Balali, G.R., S.M. Neate, E.S. Scott, D.L. Whisson and T.J. Wicks, 1995. Anastomosis group and pathogenicity of isolates of *Rhizoctonia solani* from potato crops in South Australia. *Plant Pathol.*, 44: 1050-1057.



OSCAR
PUBLISHING SERVICES