



Journal Website:
<https://theusajournals.com/index.php/ijmscr>

Copyright: Original
content from this work
may be used under the
terms of the creative
commons attributes
4.0 licence.

TO THE TREATMENT OF ACUTE BACTERIAL DESTRUCTIVE PNEUMONIA IN CHILDREN

Submission Date: May 20, 2023, Accepted Date: May 25, 2023,

Published Date: May 30, 2023

Crossref doi: <https://doi.org/10.37547/ijmscr/Volume03Issue05-10>

Mekhriddinov M.K.

Bukhara State Medical Institute Bukhara, Uzbekistan

ABSTRACT

The basis of this study was the results of treatment of 176 sick children with acute bacterial destructive pneumonia. In the treatment of this pathology, a new approach to the treatment of inflammatory infiltrates of the lung was used, based on the optimal choice of an antibiotic administered percutaneously-intrapulmonary to the lesion, which prevented the transition of the disease to a purulent-destructive stage. And also tested a differentiated method of drainage treatment of severe pleural complications of acute bacterial destructive pneumonia (ABDP) in children.

KEYWORDS

Acute bacterial destructive pneumonia, intrapulmonary administration of antibiotics, differentiated drainage treatment, empirical therapy, thoracocentesis, microflora.

INTRODUCTION

Acute bacterial destructive pneumonia (ABDP) is one of the most severe purulent-septic diseases of childhood (1,3,14,17,25). The share of various forms of destructive pneumonia accounts for 10% to 15% of the total number of pneumonias in children (2,7,19). Many scientific works have been devoted to the treatment of acute bacterial lung destruction in children, and

scientists of different generations have formed the basic principles for the diagnosis and treatment of this pathology at the present stage (4,8,11,24).

All this largely determines the level of medical care for patients with bacterial destruction of the lungs in children. Despite the success achieved in the diagnosis

and treatment of acute bacterial destructive pneumonia in children, it remains a formidable pathology in children and is one of the main causes of child mortality, which forces scientists to continue studying the features of this pathology (13,18,21).

In the list of necessary complex therapy for acute destructive pneumonia, a rather wide range of surgical methods of treatment occupies a special place, in the choice of which the form of acute destructive pneumonia plays a leading role (5,10,12,16,23).

Despite significant advances in the diagnosis and treatment of ABDP, it remains a formidable disease in young children (2,6,7,22). Over the past decade, there has been a gradual displacement of staphylococcus from etiological agents due to a wide range of gram-negative flora, as well as its association. (1,3,9,11,19). Achievements in surgical and medical treatment of ABDP have had some success, but they need to be further improved. There is a need to develop and put into practice more gentle and effective methods of drainage treatment of ABDP in childhood (5,6,7,15,20).

Aim: surgical treatment are timely and adequate sanitation of the pleural cavity and persistent re-expansion of the lung.

MATERIAL AND METHODS

The present work is based on the data of examination and treatment of 176 patients with ABDP. When

studying the clinical and radiological picture of the disease, in each case we identified several clinical groups. Most of the patients were children with exudative pleurisy - 31.3% and pyopneumothorax - 31.8%. The frequency of pleural complications was significantly higher in children under 3 years of age - 78.4%. The main contingent of patients were children of the first three years of life, including: 30.7% - under the age of one year, from one to three years old made up the majority (47.7%) of the observed patients. Among the examined patients, there were slightly more boys (54.3%) than girls (45.7%). It should be noted that in 83.7% of patients, acute bacterial destructive pneumonia occurred against the background of various concomitant diseases and complications, which were predominantly observed in children of the first three years of life. In infancy, frequent concomitant diseases were anemia (90.3%), rickets (47.3%), malnutrition (30.2%), exudative diathesis (8.9%), etc. It is important to emphasize that 94% of children history of acute viral infection and pneumonia. To solve the tasks set by us, clinical, radiological and laboratory research methods were applied.

RESULTS

When studying the etiological structure of ABDP obtained during bacteriological examination, we used two indicators: the frequency of detection of the pathogen from among all examined and from among the positive results of the tank. sowing.

Staphylococcus aureus was sown most often - in 49 cases out of 152 studies, which amounted to 32.2%. This figure rises to 62% of positive samples (79). Pseudomonas aeruginosa is in second place - 9.8% and 18%, respectively. Associations of bacteria, more often staphylococcus with Pseudomonas aeruginosa and Escherichia coli, were noted.

Thus, the traditional methods of bacteriological study of the focus of destruction, due to the low frequency of registration of pathogens and the duration of the study, do not allow the initiation of targeted, timely antibiotic therapy. In addition, it is often impossible to obtain material for a tank. sowing from the focus of the pathological process (inflammatory infiltrate, pneumothorax, bullae, fibrinous pleurisy). Considering that antibiotic therapy is urgent, it would be rational to prescribe antibiotics targeted, taking into account the pathogen and its sensitivity to antibiotics. However, the fulfillment of this requirement at the initial stages of the disease is not realistic, therefore, at the initial stages, we prescribed antibiotics based on staphylococcal etiology, either gram-negative, less often mixed. For treatment ABDP used antibiotics that have an antistaphylococcal effect and broad-spectrum antibiotics - 3-4 generation cephalosporins (ceftriaxone, rocefin, cefepime, etc.), which clinically gave a positive result from the first days of treatment. As the results of bacteriological studies were obtained, antibiotics were prescribed taking into account the

sensitivity of the isolated pathogens of the pathological process. Decrease in intoxication, normalization of temperature, appetite, decrease in the amount of contents from the drainage tube and during puncture, improvement in breathing, negative results of bacteriological studies indicated the beginning of remission. A quick identification of the causative agent of the disease ABDP allows you to choose an antibiotic that is sensitive to this type of bacteria and immediately begin antibacterial treatment of patients with ABDP.

The choice of antibiotics allows immediate initiation of antibiotic therapy by injection into the lesion. We used intrapulmonary administration of antibiotics in the infiltrate zone in 31 children. We chose the antibiotic after determining the causative agent of the disease and carried out targeted antibiotic therapy. With lobar infiltrate, the use of intrapulmonary antibiotics in the complex of therapeutic measures made it possible to achieve recovery in 93.5% of patients (29 patients). 2 patients (5%) had pneumothorax. Intrapulmonary administration of antibiotics makes it possible to create a high concentration of antibiotics in the area of inflammation, prevent the development of severe pleural complications, and in some cases contribute to the abortive course of the pathological process. In general, the method of intrapulmonary administration of antibiotics, used according to indications,

In pulmonary-pleural forms of ABDP, the main goal of drainage treatment is to remove air and fluid from the pleural cavity and straighten the lung to prevent its re-collapse. Thoracocentesis with drainage of the pleural cavity was performed in 72 patients. In the present study, according to the method of drainage treatment, patients were divided into two groups in order to determine the comparative effectiveness of the new and traditional approaches to drainage treatment.

Traditional drainage, including trocar-drainage, is usually introduced urgently, without taking into account the localization of the lesion; as a rule, along the VI-VII intercostal space, mid-axillary line, and the differentiation approach involved the introduction of a trocar-drainage at the level of the lesion, taking into account its localization. Patients who underwent thoracocentesis with drainage of the pleural cavity using a trocar-drainage according to the traditional method comprised group I (17 children). Pyopneumothorax and pneumothorax were observed in 15 patients (88%). Pyothorax in 2 patients (12%). When treated with trocar drainage, the condition of patients improved faster and the lung expanded, which was observed in 6 patients (35%). However, in 9 children (53%), a favorable clinical course was not accompanied by lung expansion. They noted a gradual, slow expansion of it. In 7 patients, the lung expanded slowly, after 2-3 weeks. In general, the use of trocar drainage made it possible to achieve recovery in 16

patients (94%). 1 patient (5%) from this group died. The length of stay of patients in the hospital was 34.1 ± 0.8 bed-days. The thoracocentesis opening on the chest wall healed 27.2 ± 1.1 days after removal of the drain.

Patients of group II - 55 children, we used the method of differentiated drainage of the pleural cavity. Pyopneumothorax was noted in 48 children (87%), pyothorax in 7 children (13%). When treated in a differentiated way, a rapid improvement in the condition with the expansion of the lung in the first hours and days was observed in 47 patients (87%), in 6 patients (11%) after a few days, only in 1 child (2%) the lung expanded in the late stages of treatment. , after the imposition of the second drainage 1-2 intercostal space above the injection site in the first drainage. The length of stay of patients in the hospital was 25.2 ± 0.5 bed-days. The thoracocentesis wound on the chest wall healed in 18.2 ± 6 days. Died 2 children (3.6%) of 55 patients in this group. The cause of adverse events here cannot be associated with the use of this method (in all the deceased, the lung remained straightened until the onset of death. In all cases there was a progressive septicopyemia).

CONCLUSIONS

1. A new approach to the treatment of inflammatory infiltrates of the lung was applied, based on the optimal choice of an antibiotic administered percutaneously-intrapulmonary to the lesion, which

prevented the transition of the disease to a purulent-destructive stage in 93% of patients.

2. A new differentiated method of drainage treatment of severe pleural complications of acute destructive pneumonia in children has been tested and introduced into clinical practice, which makes it possible to obtain a more pronounced clinical effect compared to traditional treatment.

REFERENCES

1. A.T. Akhmedov, & L.A. Narziev. (2023). PREDICTORS OF CARDIOVASCULAR COMPLICATIONS IN PATIENTS WITH CHRONIC ISCHEMIC HEART DISEASE AFTER BYPASS SURGERY. International Journal of Medical Sciences And Clinical Research, 3(03), 68–77. <https://doi.org/10.37547/ijmscr/Volume03Issue03-10>
2. A.T. Akhmedov, & Zh.R. Sharipov. (2023). THE ROLE OF CYTOKINES IN THE DEVELOPMENT OF ARTERIAL HYPERTENSION. International Journal of Medical Sciences And Clinical Research, 3(03), 59–67. <https://doi.org/10.37547/ijmscr/Volume03Issue03-09>
3. Ahmedov, A., & Kamolov, F. (2023). ECHOCARDOGRAPHIC CHANGES IN PATIENTS WITH A SURVEY OF PNEUMONIA ASSOCIATED WITH CORONAVIRUS INFECTION COVID-19. International Bulletin of Medical Sciences and Clinical Research, 3(3), 110–115. Retrieved from <https://researchcitations.com/index.php/ibmscr/article/view/883>
4. Akhmedov A.T. (2022). COMPARATIVE EVALUATION OF IMMUNOLOGICAL PARAMETERS OF LABORATORY ANIMALS WITH THYMUS AUTOIMPLANTATION IN THE DYNAMICS OF OBSERVATION. International Journal of Medical Sciences And Clinical Research, 2(11), 12–18. <https://doi.org/10.37547/ijmscr/Volume02Issue11-03>
5. An early thoracoscope approach in necrotizing pneumonia in children: a report of three cases / N. Kalfa, H. Allal, M. Lopez et al. // J. Laparoendosc. Adv. Surg. Tech. A.-2005.-Vol. 15, No. 1.-P. 18-22.
6. Barova, N. K., Tarakanov, V. A., Tsipris, A. A., Stepkin, M. A., Ubilava, I. A., Gorkova, I. S., ... & Kulish, T. A. (2013). Modern medical technologies in the treatment of acute destructive pneumonia in children. Kuban Scientific Medical Bulletin, (7), 58-59.
7. Bauer, TT Comparison of systemic cytokine levels in patients with acute respiratory distress syndrome, severe pneumonia, and controls / TT Bauer, C. Monton, A. Torres//Thorax. 2000. - Vol. 55, No. 1.-p. 1138-1145.

8. Beloborodova, N. V., Tarakanov, V. A., & Barova, N. K. (2016). Analysis of the causes of destructive pneumonia in children and the possibility of optimizing antimicrobial therapy. *Pediatrics. Journal them. GN Speransky*, 95(2), 66-71.
9. Clinical experience of managing empyema thoracic in children / FL Huang, PY Chen, JS Ma et al. // *J. Microbiol. Immunol. Infect.* 2002. - Vol. 35, No. 2. -P. 115-120.
10. Danilova, V. V., Bevez, S. I., Ovcharenko, S. S., Gurina, I. V., & Grigorova, M. V. (2018). To the question of the treatment of destructive pneumonia in children (Doctoral dissertation).
11. Diagnostic efficacy and prognostic value of serum procalcitonin concentration in patients with suspected sepsis / MJ Ruiz-Alvarez, S. Garc-a-Valdecasas, R. De Pablo et al. // *J. Intens. Care Med.* 2009. - Vol. 24, No. 1. - P. 63-71.
12. Dobrovolsky, S.R. Diagnosis and treatment of exudative pleurisy / S.R. Dobrovolsky, A.V. Belostotsky // *Surgery.* 2002. - No. 3. - S.52-57.
13. F.S., R. (2023). Morphological Features of the Ceca in the Area of Sphincters in White Laboratory Rats in Early Postnatal Ontogenesis. *Scholastic: Journal of Natural and Medical Education*, 2(4), 142–145.
14. F.S., R. (2023). Morphological Features of the Ceca in the Area of Sphincters in White Laboratory Rats in Early Postnatal Ontogenesis. *Scholastic: Journal of Natural and Medical Education*, 2(4), 142–145. Retrieved from <http://univerpubl.com/index.php/scholastic/article/view/1124>
15. Gisak S. N., Chernykh A. V., Vecherkin V. A. et al. / Modern changes in the pathomorphosis of pathogens of acute purulent destruction of the lungs and pleura in children. *Proceedings of the Russian Symposium "Ppurulent-inflammatory diseases of the lungs and pleura in children"* Krasnodar-Sochi. 2015 -p.5.
16. Isakov Yu.F., Stepanov E.A., Razumovsky A.Yu. /New technologies in pediatric surgery// *Modern technologies in pediatrics and pediatric surgery: mater. IV Congr. M., 2005.* - S. 334.
17. Kobilov, E. E., Raupov, F. S., & Mekhriddinov, M. K. (2020). A modern approach to the treatment of acute bacterial lung destruction in children. *A new day in medicine*, (4), 312-315.
18. Mashkov, A. E., Shcherbina, V. I., Tsuman, V. G., Druzyuk, E. Z., Surovikina, M. S., Nalivkt, A. E., ... & Slesarev, V. V. (2005). Modern aspects of the treatment of complicated forms of acute destructive pneumonia in children. *Almanac of Clinical Medicine*, (8-1), 240-249.
19. Mekhriddinov, M. K. (2022). Modern view on the problems of etiopathogenesis and treatment of destructive forms of pneumonia

- in children. Central Asian Journal of Medical and Natural Science, 3(2), 248-255.
20. Mekhriddinov, M. K. (2022). The role of immunological factors in the pathogenesis of the development of destructive forms of pneumonia in children. Central Asian Journal of Medical and Natural Science, 3(2), 234-238.
21. Pogorelov M. V., Danilovskikh D. A., Abushkin I. A. et al. / Analysis of antibiotic therapy in children with purulent-inflammatory diseases of the lungs and pleura at the stages of treatment. //Materials of the Russian Symposium "Purulent-inflammatory diseases of the lungs and pleura in children" Krasnodar-Sochi. 2015 -p.15.
22. Raupov, F. S., & Kobilov, E. E. EVALUATION OF THE EFFECTIVENESS OF OZONE THERAPY FOR PURULENT SURGICAL INFECTION IN CHILDREN13. LBC 57.33: 54.5 C 56, 77.
23. Raupov, F. S., & Mekhriddinov, M. K. (2021). Results of the Complex Treatment of Acute Bacterial Destruction of the Lungs in Children. Central Asian Journal of Medical and Natural Science, 146-149.
<https://doi.org/10.47494/cajmns.vio.366>
24. Raupov, F.S. (2020). Possible dysfunctions of the large intestine after resection in children. Problems of Biology and Medicine, (3), 119.
25. Sap N. A., Olenina N. V., Nekrasova E. G., Alexandrova N. S. / Thoracoscopy in acute purulent-destructive pneumonia in children. //Materials of the Russian Symposium "Ppurulent-inflammatory diseases of the lungs and pleura in children" Krasnodar-Sochi. 2015 -p.22.
26. Savenkova, M. S., Vrublevsky, S. G., Koltunov, I. E., Persianinova, E. S., Semenova, L. P., Anin, A. N., ... & Kurenkov, I. V. (2015). Destructive pneumonia in children: complex issues of diagnosis and choice of individual etiopathogenetic therapy. Farmateka, 14, 34-40.
27. Джалилова, З. О., Хасанов, К. А., & Султонов, А. А. (2021). Роль научного управления в процессе обучения высококвалифицированных врачей в новом Узбекистане. Молодой ученый, (26), 377-379.
28. Sh, Navruzova, and A.T. Akhmedov. "Autoimplantation Of Thymus In Surgical Correction Of Congenital Heart Defects." Central Asian Journal Of Medical And Natural Sciences 2.3 (2021): 88-98.
29. Shamsiev, A.M. Acute destructive pneumonia in children / AM Shamsiev. -Tashkent, 1996.-215 p.
30. Obidovna, D. Z., & Sulaymonovich, D. S. (2022). Physical activity and its impact on human health and longevity. Достижения науки и образования, (2 (82)), 120-126.

31. Obidovna, D. Z., & Sulaymonovich, D. S. (2022). THE CONCEPT OF" HEALTHY LIFESTYLE" IN PSYCHOLOGICAL RESEARCH. ResearchJet Journal of Analysis and Inventions, 3(06), 53-64.
32. Vecherkin, V. A., Toma, D. A., Ptitsyn, V. A., & Koryashkin, P. V. (2019). Destructive pneumonia in children. Russian Bulletin of Pediatric Surgery, Anesthesiology and Resuscitation, 9(4), 108-115.
33. Yusupov, Sh., Pulatov, P., Bazarov, B., & Zhaniev, H. (2015). Diagnosis and treatment of acute destructive pneumonia in children. Journal of Problems of Biology and Medicine, (1 (82)), 124-127.
34. Ахмедов А.Т., Особенности иммунной системы при врожденных пороках сердца, // EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE. – 2022. Vol. 2 No. 3 EJ MMP. – С. 35-40. [Akhmedov A. T., Features of the immune system in congenital heart defects, // EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE. – 2022. Vol. 2 no. 3 EJ MMP. – pp. 35-40 [in Russian]
35. Ахмедов А.Т., Сравнительная оценка иммунологических параметров лабораторных животных аутоимплантации тимуса в динамике наблюдения, // EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE. – 2022. - Vol. 2 No. 3: EJ MMP. С. 40-45 [Akhmedov A. T., Comparative evaluation of the immunological parameters of laboratory animals after thymus autoimplantation in the dynamics of observation, // EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE. - 2022. - Vol. 2 no. 3: EJ MMP. С. 40-45 in Russian]
36. Хидоятов, Б. А. Микроциркуляторное русло кишечника и поджелудочной железы и его особенности при экспериментальном сахарном диабете / Б. А. Хидоятов, А. Т. Ахмедов // Морфология. – 2008. – Т. 133. – № 2. – С. 145-146. – EDN JUTXTT.7. Otto CM, Pearlman AS Textbook of clinical echocardiography. Philadelphia: WB Saunders Co.; 1995:418.
37. Obidovna, D. Z., & Sulaimonovich, D. S. (2023). Influence of the Mode of Work and Recreation of the Student's Health. INTERNATIONAL JOURNAL OF HEALTH SYSTEMS AND MEDICAL SCIENCES, 2(3), 3-5.
38. Obidovna, D. Z., & Sulaymonovich, D. S. (2023). Forming a Healthy Lifestyle for Students on the Example of the Volleyball Section in Universities. EUROPEAN JOURNAL OF INNOVATION IN NONFORMAL EDUCATION, 3(3), 22-25.